

Afib is a Real Burden

Only about **1/3** of Afib is treated during heart surgery.^{1,2}

More than **33 million** people currently suffer from Afib worldwide.³

In the U.S. alone, the prevalence of Afib is projected to increase from 5.2MM in 2010 to **12.1MM** in 2032.⁴

CLINICAL BURDEN

Patients with Afib have:

- **5X** increase in stroke risk⁴
- **5X** increase in heart failure (HF) development⁵
- **46%** greater risk of death^{5,7}
- More cardiac complications⁶

PATIENT BURDEN

Patients with Afib have:

- **Decrease** in general and mental health⁸
- **Decreased** cognitive function^{9, 10, 11}
- Approximately 10 outpatient **hospital visits** and > 50 **physician encounters** per year on average¹²
- Heightened **anxiety** about medications¹³
- **Burnout** from frequent follow-up appointments¹⁴
- Up to a 47% **reduction** in quality of life¹⁵⁻¹⁹

ECONOMIC BURDEN

- Afib patients cost **\$8,700** more per year to treat²⁰
- **564,000** ED visits per year²¹
- **470,000** or **~65%** of Afib patients presenting to the emergency department (ED), are admitted each year²²
- Afib costs the U.S. health system **\$26 billion** per year²³

AFIB IS A REAL BURDEN

References:

- ¹Badhwar, V. et al. (2017). Surgical ablation of atrial fibrillation in the United States: trends and propensity matched outcomes. *Ann of Thorac Surg*, 104(2):493-500.
- ²Braid-Forbes, M.J. Health Research, 2014 CMS SAF, August 2016. NIS for volume and Medicare to look back over 3 years and obtained diagnosis.
- ³Rahman, F., Kwan, G.F., & Benjamin, E.J. (2014). Global epidemiology of atrial fibrillation. *Nat Rev Cardiol*, 11(11):639-54.
- ⁴Colilla, S. et al. (2013). Estimates of current and future incidence and prevalence of atrial fibrillation in the U.S. adult population. *Am J Cardiol*, 112(8):1142-7.
- ⁵Odutayo, A. et al. (2016). Atrial fibrillation and risks of cardiovascular disease, renal disease, and death: systematic review and meta-analysis. *BMJ*; 354:i4482.
- ⁶Fukunaga, S. et al. (2008). Effect of surgery for atrial fibrillation associated with mitral valve disease. *Ann of Thorac Surg*, 86(4):1212-7.
- ⁷Boriani, G., & Proietti, M. (2017). Atrial fibrillation prevention: an appraisal of current evidence. *Heart*, 104(11):882-7.
- ⁸Forlani, S. et al. (2006). Conversion to Sinus Rhythm by Ablation Improves Quality of Life in Patients Submitted to Mitral Valve Surgery. *Ann of Thorac Surg*, 81(3):863-7.
- ⁹Zoni-Berisso, M., Lercari, F., Carazza, T., & Domenicucci, S. (2014). Epidemiology of atrial fibrillation: European perspective. *Clin Epidemiol*, 6:213-20.
- ¹⁰Zoni-Berisso, M. et al. (2013). Frequency, patient characteristics, treatment strategies, and resource usage of atrial fibrillation (from the Italian Survey of Atrial Fibrillation Management [ISAF] study). *Am J Cardiol*, 111(5):705-11.
- ¹¹Ott, A. et al. (1997). Atrial fibrillation and dementia in a population-based study. *The Rotterdam Study. Stroke*, 28(2):316-21.
- ¹²Sullivan, E., Braithwaite, S., Dietz, K., & Hickey, C. (2010). Health services utilization and medical costs among Medicare atrial fibrillation patients. *Avalere Health*, 4(2):7.
- ¹³Wang, Y., Kong, M.C., Lee, L.H., Ng, H.J., & Ko, Y. (2014). Knowledge, satisfaction, and concerns regarding warfarin therapy and their association with warfarin adherence and anticoagulation control. *Thromb Res*, 133(4):550-4.
- ¹⁴Altioik, M., Yilmaz, M., & Rencüsogullari, I. (2015). Living with Atrial Fibrillation: An Analysis of Patients' Perspectives. *Asian Nursing Research*, 9(4):305-11.
- ¹⁵Nazli, C. et al. (2016). Impaired quality of life in patients with intermittent atrial fibrillation. *Anatol J Cardiol*, 16(4):250-5.
- ¹⁶Thrall, G., Lane, D., Carroll, D., & Lip, G.Y. (2006). Quality of life in patients with atrial fibrillation: a systematic review. *Am J Med*, 119(5):448.e1-19.
- ¹⁷Hagens, V.E. et al. (2004). Effect of rate or rhythm control on quality of life in persistent atrial fibrillation. Results from the Rate Control Versus Electrical Cardioversion (RACE) Study. *J Am Coll Cardiol*, 43(2):241-7.
- ¹⁸Hoegh, V. et al. (2016). Association between the diagnosis of atrial fibrillation and aspects of health status: a Danish cross-sectional study. *Scand J Caring Sci*, 30(3):507-17.
- ¹⁹Dorian, P. et al. (2000). The impairment of health-related quality of life in patients with intermittent atrial fibrillation: implications for the assessment of investigational therapy. *J Am Coll Cardiol*, 36(4):1303-9.
- ²⁰Centers for Disease Control and Prevention. (2017). Atrial Fibrillation Fact Sheet. Accessed Dec 2018. Retrieved from https://www.cdc.gov/dhds/data_statistics/fact_sheets/fs_atrial_fibrillation.htm.
- ²¹McDonald, A.J., Pelletier, A.J., Ellinor, P.T., & Camargo Jr, C.A. (2008). Increasing US emergency department visit rates and subsequent hospital admissions for atrial fibrillation from 1993 to 2004. *Ann Emerg Med*, 51(1):58-65.
- ²²Rozen, G. et al. (2018). Emergency Department Visits for Atrial Fibrillation in the United States: Trends in Admission Rates and Economic Burden From 2007 to 2014. *J Am Heart Assoc*, 7(15), pii:e009024.
- ²³Kim, M.H., Johnston, S.S., Chu, B.C., Dalal, M.R., & Schulman, K.L. (2011). Estimation of total incremental health care costs in patients with atrial fibrillation in the United States. *Circ Cardiovasc Qual Outcomes*, 4(3):313-20.