


Surgical Ablation Guidelines are Advancing



COR = Class of Recommendation
 LOE = Level of Evidence

Sources: + Article in Press. [https://www.heartrhythmjournal.com/article/S1547-5271\(24\)00261-3/fulltext](https://www.heartrhythmjournal.com/article/S1547-5271(24)00261-3/fulltext) (accessed 4/10/2024). Heart Rhythm Society, the European Society of Cardiology, the Asia Pacific Heart Rhythm Society, and the Latin American Heart Rhythm Society 2024. + Hybrid ablation type of evidence META (meta-analysis); LAAE type of evidence RAND (randomized controlled); nomenclature did not use LOE classification. ^Advice TO DO/RAND. †Advice TO DO/META.

Wyler von Ballmoos, M.C. et al. (2024). The Society of Thoracic Surgeons 2023 Clinical Practice Guidelines for the Surgical Treatment of Atrial Fibrillation.

Members, W. C., et al. (2023). 2023 ACC/AHA/ACCP/HRS Guideline for the Diagnosis and Management of Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. Journal of the American College of Cardiology.

January, C. T., et al. (2019). 2019 AHA/ACC/HRS Focused Update of the 2014 AHA/ACC/HRS Guideline for the Management of Patients With Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. Circulation, CIR-0000000000000665.

Badhwar, et al. (2017). The Society of Thoracic Surgeons 2017 Clinical Practice Guidelines for the Surgical Treatment of Atrial Fibrillation. Ann Thorac Surg, 103(1):329-41. †MVR LOE A; AVR,CABG LOE B.

January, C.T., et al. (2014). 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. J Am Coll Cardiol, 64(21):e1-76.

*Calkins, H., et al. (2017). 2017 HRS/EHRA/ECAS/APHRS/SOLAECE expert consensus statement on catheter and surgical ablation of atrial fibrillation. Heart Rhythm, 14(10):e275-444. AVR/CABG concomitant ablation Class I LOR for symptomatic persistent and long-standing persistent “refractory or intolerant to at least one Class 1 or 3 antiarrhythmic medication.”

Meier, B., et al. (2014). EHRA/EAPCI expert consensus statement on catheter-based left atrial appendage occlusion. Europace, 16(10):1397-416.

Cox, J.L., et al. (1991). Dr. Cox performed first surgical ablation using maze I; Successful surgical treatment of atrial fibrillation. Review and clinical update. JAMA, 266 (14):1976-80.

Isabelle C Van Gelder, Michiel Rienstra, Karina V Bunting, Ruben Casado-Arroyo, Valeria Caso, Harry J G M Crijns, Tom J R De Potter, Jeremy Dwight, Luigina Guasti, Thorsten Hanke, et al. (2024). 2024 ESC Guidelines for the management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): Developed by the task force for the management of atrial fibrillation of the European Society of Cardiology (ESC), with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. Endorsed by the European Stroke Organisation (ESO), European Heart Journal, ehae176.